

Hip Pain in Adults: Evaluation and Differential Diagnosis

Rachel Chamberlain, MD, CAQSM, University of New Mexico, Albuquerque, New Mexico

Adults commonly present to their family physicians with hip pain, and diagnosing the cause is important for prescribing effective therapy. Hip pain is usually located anteriorly, laterally, or posteriorly. Anterior hip pain includes referred pain from intra-abdominal or intrapelvic causes; extra-articular etiologies, such as hip flexor injuries; and intra-articular etiologies. Intra-articular pain is often caused by a labral tear or femoroacetabular impingement in younger adults or osteoarthritis in older adults. Lateral hip pain is most commonly caused by greater trochanteric pain syndrome, which includes gluteus medius tendinopathy or tear, bursitis, and iliotibial band friction. Posterior hip pain includes referred pain such as lumbar spinal pathology, deep gluteal syndrome with sciatic nerve entrapment, ischiofemoral impingement, and hamstring tendinopathy. In addition to the history and physical examination, radiography, ultrasonography, or magnetic resonance imaging may be needed for a definitive diagnosis. Radiography of the hip and pelvis should be the initial imaging test. Ultrasound-guided anesthetic injections can aid in the diagnosis of an intra-articular cause of pain. Because femoroacetabular impingement, labral tears, and gluteus medius tendon tears typically have good surgical outcomes, advanced imaging and/or early referral may improve patient outcomes. (*Am Fam Physician*. 2021;103(2):81-89. Copyright © 2021 American Academy of Family Physicians.)

Hip pain is common in adults of all ages and activity levels. In nonelite adult soccer players, hip and groin injuries represent 28% to 45% of all injuries in women and 49% to 55% in men.¹ The prevalence of the cam deformity (deformity of the femoral head) is 41% in nonelite male soccer players and 17% in male nonathletes.² In adults older than 45 years, 6.7% to 9.7% have osteoarthritis of the hip, and one in four adults will develop symptomatic hip osteoarthritis in their lifetime.³ In the United States in 2009, hip replacements accounted for \$13.7 billion in health care costs.³

Approach to Evaluation

Hip pain is often localized to one of three locations: anterior, lateral, or posterior (*Figure 1*). A focused history and physical examination can help differentiate the causes of hip pain

(*Table 1*).²⁻²⁰ Diagnosing the cause of hip pain is important for prescribing effective therapy.

The history should include personal history of developmental hip dysplasia, slipped capital femoral epiphysis, sports activities, and injuries; family history of hip problems; and the location and quality of pain, aggravating and alleviating factors, and mechanical symptoms.^{4,8} Physical examination should include gait analysis with particular attention to antalgic or Trendelenburg gait, evaluation of the range of motion in the hip joint and associated pain, strength testing of the muscles overlying the hip joint, palpation of the painful area, and special tests as indicated.

If imaging is performed in the evaluation of a patient with undifferentiated chronic hip pain, standing anteroposterior hip and pelvic radiography is typically the initial imaging study.^{4,21} Magnetic resonance imaging (MRI) or ultrasonography may be helpful in the diagnosis, depending on history and physical examination findings.²¹⁻²³

Anterior Hip Pain

Intra-articular hip pain predominately presents anteriorly.^{4,5} In young adults, anterior hip or groin pain that is aggravated by hip flexion or rotation

CME This clinical content conforms to AAFP criteria for CME. See CME Quiz on page 79.

Author disclosure: No relevant financial affiliations.


Patient information: A handout on this topic is available at <https://www.aafp.org/aafp/2021/0101/p81-s1.html>.

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	Comments
If imaging is performed in the evaluation of a patient with undifferentiated chronic hip pain, standing anteroposterior hip and pelvic radiographs should be the first choice. ^{4,21}	C	Expert opinion and consensus guidelines
For patients with anterior hip pain and history suggestive of a labral tear, stress fracture of the femoral neck, or early avascular necrosis, magnetic resonance imaging should be performed for accurate diagnosis. ^{5,11,12,21-23}	C	Expert opinion and reviews of prospective and randomized trials
For intra-articular pain, ultrasound-guided anesthetic injection of the hip may be diagnostic, and corticosteroid injection may be therapeutic. ³⁰	C	Clinical review and expert opinion
For patients with greater trochanteric pain syndrome not responding to conservative therapy, ultrasonography or magnetic resonance imaging should be considered to evaluate for gluteus medius tendon tears. ^{15,16,21}	C	Clinical reviews and expert opinion

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to [www.aafp.org/sort](#).

FIGURE 1



Localization of hip pain. (A) Anterior view. (B) Posterior view.

Illustration by Todd Buck

Reprinted with permission from Wilson JJ, Furukawa M. Evaluation of the patient with hip pain. *Am Fam Physician*. 2014;89(1):27-34.

warrants evaluation for intra-articular pathologies. Hip flexor strains, tears, and avulsion fractures can cause anterior hip pain, with the patient history often including a sports-related or traumatic incident consistent with a flexor injury.⁵ Because referred pain from intra-abdominal

problems can present as anterior hip pain, the abdomen should be examined for gastrointestinal causes of pain, such as a mass; appendicitis; hernia; or pain originating in the bladder (e.g., from a mass) or the female reproductive system (e.g., from ovarian cysts).^{4,6}

TABLE 1

Differential Diagnosis of Hip Pain

Type of pain	History	Physical examination
Anterior		
Referred		
Intra-abdominal or intrapelvic ⁴⁻⁶	Pain associated with urinary or bowel symptoms, cyclic pain associated with menses	Abdominal and/or pelvic examination
Extra-articular		
Flexor tendon ^{5,6}	Overuse activities, acute strain or injury with hip flexion activities	Pain over the hip bony prominence, anterior superior iliac spine, anterior inferior iliac spine, or pubic symphysis; pain with hip flexion strength testing
Intra-articular		
Femoroacetabular impingement ^{2,5,7,8}	Young, athletic patient; gradual onset; pain with hip range of motion; history of slipped capital femoral epiphysis or developmental dysplasia	Positive FADDIR and FABER test results
Labral tear ^{5,9}	Young, athletic patient; acute injury (vs. gradual onset); pain with hip range of motion; mechanical symptoms	Positive FADDIR and FABER test results
Femoral neck stress fracture ^{5,10}	Overuse/overtraining, energy imbalance in athletes	Antalgic gait, pain with range of motion and ambulating
Avascular necrosis ^{11,12}	Middle or older age, smoking, alcohol use, systemic corticosteroid use, hemoglobinopathies, chemotherapy, metabolic syndrome, and obesity	Antalgic gait, pain with range of motion, limited range of motion
Osteoarthritis ^{3,4,7,13}	Older age, gradual onset, pain with sitting or ambulating for long periods	Antalgic gait, pain with flexion and internal and external rotation, limited range of motion
Hip fracture ^{4,14}	Older age, osteoporosis, fall/trauma	Inability to walk on the affected limb; shortened, externally rotated, abducted leg
Lateral		
Greater trochanteric pain syndrome, including bursitis, gluteus medius tendinopathy or tear, external snapping, or iliotibial band friction ^{7,15,16}	No injury, middle age, female sex, overweight, pain with sleeping on affected hip, pain aggravated by physical activity or sitting for long periods	Tenderness to palpation over the lateral hip/greater trochanter, Trendelenburg gait or positive Trendelenburg test, positive resisted external derotation test
Posterior		
Referred pain		
Intra-abdominal or intrapelvic ^{4,17}	Pain associated with urinary or bowel symptoms, cyclic pain associated with menses	Abdominal and/or pelvic examination
Deep gluteal syndrome ^{17,18}	Deep buttock pain; no injury; worse with sitting, especially in a car; sciatica (burning pain shooting down the leg)	Seated piriformis stretch test
Ischiofemoral impingement ¹⁹	Gradual onset of deep buttock pain that worsens with activities requiring a long stride (e.g., running)	Long-stride walking test
Lumbar spine or muscle ^{4,17}	Pain in the low back (above L5) and hip/buttock, history of lumbar spinal problems	Tenderness over the lumbar spine or lumbar musculature above L5
Sacroiliac joint pain ¹⁷	No history of lumbar spinal issues	Tenderness over the sacroiliac joint, no tenderness above L5
Proximal hamstring tendinopathy or tear ²⁰	Overuse injury with hip extension activities (vs. acute injury with forceful hip extension)	Tenderness to palpation over the ischial tuberosity, pain with hamstring strength testing; acute tears cause ecchymosis of the posterior thigh

FABER = flexion abduction external rotation; FADDIR = flexion adduction internal rotation.

Information from references 2-20.

FEMOROACETABULAR IMPINGEMENT

Femoroacetabular impingement is one of the most common causes of hip pain in young adults.²⁴ It can be caused by a cam deformity, which is bony overgrowth of the femoral head and neck, a pincer deformity of the acetabulum (too much coverage of the femoral head), or both (Figure 2²⁵). Femoroacetabular impingement often has a gradual onset without a specific injury. It usually presents earlier in the disease process with less bony changes in athletes than in nonathletes and is more bothersome to athletes whose activities require hyperflexion and wide range of motion at the hip joint.^{2,5,9} Positive results on the flexion

adduction internal rotation and flexion abduction external rotation tests (Figures 3 and 4) are indicative of intra-articular hip pathology.^{4,7}

Radiography is used to help diagnose femoroacetabular impingement, although the preferred radiographic views may vary. Standing anteroposterior radiographs of the pelvis may be most helpful for diagnosing of the pincer deformity,²⁶ whereas Meyer lateral and 90-degree Dunn views may be best to visualize the cam deformity.²⁷

LABRAL TEARS

Patients with labral tears may present with anterior hip pain and a history of a sports-related or traumatic injury. They may also be associated with repetitive motions. Labral tears may cause a popping, catching, or clicking sound associated with activities such as dance, gymnastics, hockey, basketball, and soccer.^{2,5,9} Physical examination for labral tears should include flexion adduction internal rotation and flexion abduction external rotation tests.⁷


A standing radiograph should be the initial imaging test.²⁴ Previously, magnetic resonance arthrography (1.5 tesla) with gadolinium injection of the hip was the diagnostic standard for labral tears. However, with recent advances to 3-tesla MRI and specialized hip protocols, noncontrast 3-tesla MRI is as sensitive and specific as magnetic resonance arthrography and does not require a procedure for contrast injection.^{22,23} Physicians should consult with a local radiologist to determine the most appropriate test in their area.

Labral tears and femoroacetabular impingement are often comorbid conditions in young active patients. Athletes are more likely to require surgical intervention for these conditions, especially those with both conditions.⁹

FEMORAL NECK STRESS FRACTURES

Stress fractures of the femoral neck are typically associated with overuse and may also be associated with energy imbalance in athletes. Femoral neck stress fractures are more common in women than men but should not be excluded in men with a history of overuse.¹⁰ Early in the disease process, stress fractures are typically not visible on radiographs and therefore MRI is required for definitive diagnosis. Early diagnosis of a femoral neck stress fracture is important because conversion to a complete fracture can be a devastating injury.^{5,28}

FIGURE 2



Causes of femoroacetabular impingement. A pincer deformity results from excessive overhang of the acetabulum, which causes the labrum to be impinged between the acetabulum and femoral head when the hip is flexed. A cam deformity occurs when exostosis along the femoral head and neck impinges the labrum against the acetabulum.

Illustration by Dave Klemm

Reprinted with permission from Kuhlman GS, Domb BG. Hip impingement: identifying and treating a common cause of hip pain. *Am Fam Physician.* 2009;80(12):1431.

FIGURE 3



Flexion adduction internal rotation test. The examiner (A) passively flexes then (B) adducts and internally rotates the hip. The result is positive if pain is reproduced in the anterior hip/groin area. This test has been shown to have a sensitivity of 59% to 100% and specificity of 4% to 75% for intra-articular hip pathology.^{4,7}

FIGURE 4



Flexion abduction external rotation test. The examiner (A) passively flexes and then (B) abducts and externally rotates the hip. The result is positive if pain is reproduced in the anterior hip/groin area. This test has been shown to have 42% to 81% sensitivity and 18% to 25% specificity for intra-articular hip pathology.^{4,7}

AVASCULAR NECROSIS

Avascular necrosis of the femoral head most commonly presents in middle-aged to older adults. Risk factors include alcohol use, smoking, systemic corticosteroid use, hemoglobinopathies, chemotherapy, metabolic syndrome, and obesity. Early detection of avascular necrosis can be joint sparing, but no physical examination finding is specific for this diagnosis.¹¹ Late-stage disease may be visible on a radiograph, but earlier diagnosis often requires MRI or computed tomography.^{12,21}

OSTEOARTHRITIS

In older adults, osteoarthritis of the femoroacetabular joint is the most common cause of anterior hip pain.³ It can lead to significant morbidity and decrease in physical activity. Osteoarthritis of the hip typically has a gradual onset, but some patients recall a specific injury or fall. Patients with this condition may have pain with sitting and ambulating for long periods and may have an antalgic gait. Physical examination maneuvers such as flexion and internal and external rotation may reproduce pain, and range of motion may be decreased.^{4,6}

Standing anteroposterior radiography of the pelvis is the radiologic test of choice and will show joint space narrowing and osteophyte formation. However, presence of these findings does not always correlate with symptom severity.^{7,13,29} Ultrasound-guided anesthetic injection of the hip joint may help differentiate an intra-articular cause of pain from other causes (e.g., lumbar spine or extra-articular pain), and corticosteroid injection may be therapeutic for intra-articular pain.³⁰

HIP FRACTURES

Hip fractures are more common in older adults and often present after a fall or other trauma or may be associated with osteoporosis. Physical examination usually reveals an inability to walk on the affected limb and a shortened, externally rotated, abducted leg while in the supine position.¹⁴ Most hip fractures are visible on a radiograph and require surgical fixation.⁴

Lateral Hip Pain

The most common cause of lateral hip pain is greater trochanteric pain syndrome (previously called greater trochanteric bursitis).¹⁵ Although

this syndrome may be associated with bursitis, gluteus medius tendinopathy or tears are now thought to be more common.^{16,31} Iliotibial band friction and external snapping hip can contribute to greater trochanteric pain syndrome.¹⁵

Greater trochanteric pain syndrome presents as lateral hip pain aggravated by ambulation or other physical activities, sitting for long periods, and sleeping on the affected hip. It most commonly affects women 40 to 60 years of age.¹⁶ There is typically no inciting injury. The patient may walk with a Trendelenburg gait or have positive findings on the Trendelenburg test or resisted external derotation test.^{7,16} (*Figure 5*).

If a patient with greater trochanteric pain syndrome does not improve with anti-inflammatory medications and physical therapy, a gluteus medius tendon tear should be considered. MRI and musculoskeletal ultrasonography performed by an experienced sonographer are sufficiently sensitive and specific for diagnosing a gluteus medius tendon tear.^{15,16,21} Referral to an orthopedic hip specialist is often indicated for large partial or complete tears because surgery is typically associated with good outcomes in patients with this condition.^{15,16}

Posterior Hip Pain

The cause of posterior hip pain can be difficult to diagnose. The differential diagnosis includes musculoskeletal causes and referred pain from intrapelvic and gynecologic issues. Patients with intrapelvic problems may have a history of cyclic pain associated with menses or urinary or bowel symptoms.¹⁷

PIRIFORMIS AND DEEP GLUTEAL SYNDROME

Piriformis syndrome is thought to be a result of the piriformis muscle entrapping the sciatic nerve, causing hip and buttock pain and sciatica (burning pain shooting down the leg). Piriformis syndrome is a subset of deep gluteal syndrome, which includes entrapment of the sciatic nerve and/or pudendal nerve by the piriformis muscle, gemelli-obturator internus, or proximal hamstrings.¹⁸ Patients with deep gluteal syndrome have deep buttock pain that is aggravated by sitting and sciatica symptoms. The seated piriformis stretch test (*Figure 6*) may reproduce this pain.¹⁸

Although conservative treatments are often helpful, MRI may be needed to identify

HIP PAIN IN ADULTS

pathology in the deep gluteal muscles or sciatic and pudendal nerves. Additionally, electrodiagnostic nerve testing can help localize the area of nerve entrapment.¹⁸

ISCHIOFEMORAL IMPINGEMENT

Ischiofemoral impingement is impingement of the quadratus femoris muscle and nerve between the proximal femur at the level of the lesser trochanter and the ischial tuberosity.¹⁹ Patients with ischiofemoral impingement have gradual onset of deep buttock pain that is worsened with activities requiring a long stride, such as running. The

long-stride walking test (*Figure 7*) is the most sensitive and specific test for this condition.^{18,19}

LUMBAR SPINE AND SACROILIAC JOINT PATHOLOGY

Lumbar spinal issues can present as posterior hip pain. Patients typically have pain in the lumbar spine or musculature and in the posterior hip/buttock area and may report previous lumbar spinal problems. Radiography of the lumbar spine may show degenerative disease, and MRI can help identify disk herniation or nerve entrapment.¹⁷

FIGURE 5



Tests for gluteal tendinopathy. (A) Modified Trendelenburg test. Patients are instructed to stand on the affected leg/hip and lift the other leg for 30 seconds. The result is positive if the iliac crest falls below the standing side, showing weakness in hip abductors (gluteus muscles). A positive finding on the Trendelenburg test has been shown to have 23% to 97% sensitivity and 77% to 96% specificity for gluteal tendinopathy.^{4,7} (B) Resisted external derotation test. While the patient lies on a table, the hip is passively flexed to 90 degrees, then externally rotated. The patient is asked to return the leg to the same axis as the table (0 degrees rotation), pushing against the examiner's (resisting) hand. The result is positive if pain is reproduced over the lateral hip. This test has been shown to have 88% sensitivity and 97% specificity for gluteal tendinopathy.⁷

FIGURE 6



Seated piriformis stretch test. The patient is seated with 90 degrees of hip flexion. The examiner extends the knee and passively moves the hip into adduction and internal rotation while palpating just lateral to the ischium. The result is positive if pain is reproduced at the piriformis muscle. This test has been shown to have a 91% sensitivity and 80% specificity for deep gluteal syndrome.¹⁸

FIGURE 7



Long-stride walking test. The patient is instructed to take a long step on the unaffected leg with the hip pointed forward. This narrows the space between the lesser trochanter and ischium on the posterior (affected) hip, which can reproduce pain caused by ischiofemoral impingement. This test has been shown to have a sensitivity of 94% and specificity of 85% for ischiofemoral impingement.^{18,19}

Sacroiliac joint dysfunction and/or arthritis may also present as posterior hip pain. The most common physical examination finding is tenderness to palpation over the sacroiliac joint. Sacroiliac pain typically does not occur above the L5 level, which would indicate a lumbar spinal etiology of pain.¹⁷ Radiography may show sacroiliac joint arthritis. Image-guided injections may be a helpful diagnostic and therapeutic tool. If diagnosis is uncertain, MRI can show inflammation or arthritis at the sacroiliac joint.¹⁷

HAMSTRING INJURIES

Posterior hip/buttock pain around the ischium may be indicative of a hamstring strain, tear, or avulsion. Patients may have a history of a traumatic, sports-related, or overuse injury.

A complete hamstring tear or avulsion often causes ecchymosis of the posterior thigh.²⁰ Hamstring tendinopathy or partial tears are typically exacerbated by hamstring strength testing but do not cause obvious bruising or deformity. Patients with acute tears should be referred to an orthopedic surgeon.²⁰ If it is unclear whether the patient has a hamstring injury, MRI may be helpful in determining the diagnosis.¹⁷

This article updates previous articles on this topic by Wilson and Furukawa,⁴ and O’Kane.⁶

Data Sources: A PubMed search was conducted using the following key words, with filters set for the previous five years and humans only: hip pain, anterior hip pain, femoroacetabular impingement, gluteus medius tendinitis, hip physical exam, 3-tesla hip MRI, ultrasound guided hip injection, greater trochanteric

pain syndrome, deep gluteal syndrome, ischiofemoral syndrome, hamstring avulsion. Additional searches included the Cochrane Database of Systematic Reviews and Essential Evidence Plus. Search dates: November 8, 2019, to September 30, 2020.

The Author

RACHEL CHAMBERLAIN, MD, CAQSM, is the associate program director of the Sports Medicine Fellowship and an assistant professor in the Department of Family and Community Medicine at the University of New Mexico, Albuquerque.

Address correspondence to Rachel Chamberlain, MD, CAQSM, 1 University of New Mexico, Albuquerque, NM 87131 (email: rchamberlain@salud.unm.edu). Reprints are not available from the author.

References

- Langhout R, Weir A, Litjes W, et al. Hip and groin injury is the most common non-time-loss injury in female amateur football. *Knee Surg Sports Traumatol Arthrosc*. 2019; 27(10):3133-3141.
- Thorborg K, Rathleff MS, Petersen P, et al. Prevalence and severity of hip and groin pain in sub-elite male football: a cross-sectional cohort study of 695 players. *Scand J Med Sci Sports*. 2017;27(1):107-114.
- Murphy L, Helmick CG. The impact of osteoarthritis in the United States: a population-health perspective. *Am J Nurs*. 2012;112(3 suppl 1):S13-S19.
- Wilson JJ, Furukawa M. Evaluation of the patient with hip pain. *Am Fam Physician*. 2014;89(1):27-34. Accessed August 19, 2020. <https://www.aafp.org/afp/2014/0101/p27.html>
- Dick AG, Houghton JM, Bankes MJK. An approach to hip pain in a young adult. *BMJ*. 2018;361:k1086.
- O'Kane JW. Anterior hip pain. *Am Fam Physician*. 1999; 60(6):1687-1696. Accessed August 19, 2020. <https://www.aafp.org/afp/1999/1015/p1687.html>
- Reiman MP, Goode AP, Hegedus EJ, et al. Diagnostic accuracy of clinical tests of the hip: a systematic review with meta-analysis. *Br J Sports Med*. 2013;47(14):893-902.
- Kamegaya M, Saisu T, Nakamura J, et al. Drehmann sign and femoro-acetabular impingement in SCFE. *J Pediatr Orthop*. 2011;31(8):853-857.
- Cianci A, Sugimoto D, Stracciolini A, et al. Nonoperative management of labral tears of the hip in adolescent athletes: description of sports participation, interventions, comorbidity, and outcomes. *Clin J Sport Med*. 2019;29(1): 24-28.
- Kupferer KR, Bush DM, Cornell JE, et al. Femoral neck stress fracture in Air Force basic trainees. *Mil Med*. 2014; 179(1):56-61.
- Lamb JN, Holton C, O'Connor P, et al. Avascular necrosis of the hip. *BMJ*. 2019;365:l2178.
- Guerado E, Caso E. The physiopathology of avascular necrosis of the femoral head: an update. *Injury*. 2016; 47(suppl 6):S16-S26.
- Beumer L, Wong J, Warden SJ, et al. Effects of exercise and manual therapy on pain associated with hip osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2016;50(8):458-463.
- LeBlanc KE, Muncie HL Jr., LeBlanc LL, et al. Hip fracture: diagnosis, treatment, and secondary prevention. *Am Fam Physician*. 2014;89(12):945-951. Accessed August 19, 2020. <https://www.aafp.org/afp/2014/0615/p945.html>
- Redmond JM, Chen AW, Domb BG. Greater trochanteric pain syndrome. *J Am Acad Orthop Surg*. 2016;24(4): 231-240.
- LaPorte C, Vasaris M, Gossett L, et al. Gluteus medius tears of the hip: a comprehensive approach. *Phys Sportsmed*. 2019;47(1):15-20.
- Gómez-Hoyos J, Martin RL, Martin HD. Current concepts review: evaluation and management of posterior hip pain. *J Am Acad Orthop Surg*. 2018;26(17):597-609.
- Park JW, Lee Y-K, Lee YJ, et al. Deep gluteal syndrome as a cause of posterior hip pain and sciatica-like pain. *Bone Joint J*. 2020;102-B(5):556-567.
- Hernando MF, Cerezal L, Pérez-Carro L, et al. Evaluation and management of ischiofemoral impingement: a patho-physiologic, radiologic, and therapeutic approach to a complex diagnosis. *Skeletal Radiol*. 2016;45(6):771-787.
- Degen RM. Proximal hamstring injuries: management of tendinopathy and avulsion injuries. *Curr Rev Musculoskelet Med*. 2019;12(2):138-146.
- American College of Radiology ACR Appropriateness Criteria. Chronic hip pain. Accessed June 5, 2020. <https://acsearch.acr.org/docs/69425/Narrative/>
- Carstensen SE, McCrum EC, Pierce JL, et al. Magnetic resonance imaging (MRI) and hip arthroscopy correlations. *Sports Med Arthrosc Rev*. 2017;25(4):199-209.
- Crespo-Rodríguez AM, De Lucas-Villarrubia JC, Pastrana-Ledesma M, et al. The diagnostic performance of non-contrast 3-tesla magnetic resonance imaging (3-T MRI) versus 1.5-tesla magnetic resonance arthrography (1.5-T MRA) in femoro-acetabular impingement. *Eur J Radiol*. 2017;88:109-116.
- Haldane CE, Ekhtiari S, de Sa D, et al. Preoperative physical examination and imaging of femoroacetabular impingement prior to hip arthroscopy—a systematic review. *J Hip Preserv Surg*. 2017;4(3):201-213.
- Kuhlman GS, Domb BG. Hip impingement: identifying and treating a common cause of hip pain. *Am Fam Physician*. 2009;80(12):1429-1434. Accessed August 19, 2020. <https://www.aafp.org/afp/2009/1215/p1429.html>
- Jackson TJ, Estess AA, Adamson GJ. Supine and standing AP pelvis radiographs in the evaluation of pincer femoroacetabular impingement. *Clin Orthop Relat Res*. 2016; 474(7):1692-1696.
- Atkins PR, Shin Y, Agrawal P, et al. Which two-dimensional radiographic measurements of cam femoroacetabular impingement best describe the three-dimensional shape of the proximal femur? *Clin Orthop Relat Res*. 2019;477(1): 242-253.
- Adkins SB III, Figler RA. Hip pain in athletes. *Am Fam Physician*. 2000;61(7):2109-2118. Accessed August 19, 2020. <https://www.aafp.org/afp/2000/0401/p2109.html>
- Kim C, Nevitt MC, Niu J, et al. Association of hip pain with radiographic evidence of hip osteoarthritis: diagnostic test study. *BMJ*. 2015;351:h5983.
- Lynch TS, Oshlag BL, Bottiglieri TS, et al. Ultrasound-guided hip injections. *J Am Acad Orthop Surg*. 2019;27(10):e451-e461.
- Brennan KL, Allen BC, Maldonado YM. Dry needling versus cortisone injection in the treatment of greater trochanteric pain syndrome: a noninferiority randomized clinical trial. *J Orthop Sports Phys Ther*. 2017;47(4):232-239.